arXiv:1109.4122v1 [math.CO] 19 Sep 2011

Pairs of heavy subgraphs for Hamiltonicity of 2-connected graphs*

Binlong Li[a], Zdeněk Ryjáček[b,†], Ying Wang[a] and Shenggui Zhang[a,‡]

[a] Department of Applied Mathematics, Northwestern Polytechnical University,
Xi'an, Shaanxi 710072, P.R. China

[b] Department of Mathematics, University of West Bohemia and
Institute for Theoretical Computer Science, Charles University,
30614 Pilsen, Czech Republic

Abstract

Let G be a graph on n vertices. An induced subgraph H of G is called heavy if there exist two nonadjacent vertices in H with degree sum at least n in G. We say that G is H-heavy if every induced subgraph of G isomorphic to H is heavy. For a family $\mathcal{H}$ of graphs, G is called $\mathcal{H}$-heavy if G is H-heavy for every $H\in\mathcal{H}$. In this paper we characterize all connected graphs R and S other than P_3 (the path on three vertices) such that every 2-connected $\{R,S\}$-heavy graph is Hamiltonian. This extends several previous results on forbidden subgraph conditions for Hamiltonian graphs.

Keywords: Forbidden subgraph; Heavy subgraph; Hamilton cycle

1 Introduction

We use Bondy and Murty [2] for terminology and notation not defined here and consider finite simple graphs only.

Let G be a graph. For a vertex $v\in V(G)$ and a subgraph H of G, we use $N_H(v)$ to denote the set, and $d_H(v)$ the number, of neighbors of v in H, respectively. We call $d_H(v)$ the *degree* of v in H. For $x,y\in V(G)$, an (x,y)-*path* is a path P connecting x and y; the vertex x will be called the *origin* and y the *terminus* of P. For $X,Y\subset V(G)$, an (X,Y)-*path* is a path having its origin in X and terminus in Y. If $x,y\in V(H)$, the

---

[*] This paper was supported by NSFC (No. 10871158).

[†] Research partially supported by grant 1M0545 of the Czech Ministry of Education

[‡] Corresponding author. E-mail address: sgzhang@nwpu.edu.cn (S. Zhang).

distance between x and y in H, denoted $d_H(x,y)$, is the length of a shortest (x,y)-path in H. When no confusion occurs, we will denote $N_G(v)$, $d_G(v)$ and $d_G(x,y)$ by $N(v)$, $d(v)$ and $d(x,y)$, respectively.

Let G be a graph on n vertices. If a subgraph G' of G contains all edges $xy \in E(G)$ with $x, y \in V(G')$, then G' is called an *induced subgraph* of G. For a given graph H, we say that G is *H-free* if G does not contain an induced subgraph isomorphic to H. For a family $\mathcal{H}$ of graphs, G is called $\mathcal{H}$-free if G is H-free for every $H \in \mathcal{H}$. If H is an induced subgraph of G, we say that H is *heavy* if there are two nonadjacent vertices in $V(H)$ with degree sum at least n in G. The graph G is called *H-heavy* if every induced subgraph of G isomorphic to H is heavy. For a family $\mathcal{H}$ of graphs, G is called *$\mathcal{H}$-heavy* if G is H-heavy for every $H \in \mathcal{H}$. Note that an H-free graph is also H-heavy.

The graph $K_{1,3}$ is called the *claw*, its (only) vertex of degree 3 is called its *center* and the other vertices are the *end vertices*. In this paper, instead of $K_{1,3}$-free ($K_{1,3}$-heavy), we use the terminology claw-free (claw-heavy).

The following characterization of pairs of forbidden subgraphs for the existence of Hamilton cycles in graphs is well known.

Theorem 1 (Bedrossian [1]). *Let R and S be connected graphs with $R, S \neq P_3$ and let G be a 2-connected graph. Then G being $\{R, S\}$-free implies G is Hamiltonian if and only if (up to symmetry) $R = K_{1,3}$ and $S = P_4, P_5, P_6, C_3, Z_1, Z_2, B, N$ or W (see Fig. 1).*



Fig. 1. Graphs P_i, C_3, Z_i, B, N and W.

Our aim in this paper is to consider the corresponding heavy subgraph condition for a graph to be Hamiltonian. First, we notice that every 2-connected P_3-heavy graph contains a Hamilton cycle. This can be easily deduced from the following result.

Theorem 2 (Fan [5]). *Let G be a 2-connected graph. If* $\max\{d(u), d(v)\} \geq n/2$ *for every pair of vertices with distance 2 in G, then G is Hamiltonian.*

It is not difficult to see that P_3 is the only connected graph S such that every 2-connected S-heavy graph is Hamiltonian. So we have the following problem.

Problem 1. Which two connected graphs R and S other than P_3 imply that every 2-connected $\{R, S\}$-heavy graph is Hamiltonian?

By Theorem 1, we get that (up to symmetry) $R = K_{1,3}$ and S must be some of the graphs P_4, P_5, P_6, C_3, Z_1, Z_2, B, N or W.

In this paper we prove the following results.

Theorem 3. *If G is a 2-connected $\{K_{1,3}, W\}$-heavy graph, then G is Hamiltonian.*

Theorem 4. *If G is a 2-connected $\{K_{1,3}, N\}$-heavy graph, then G is Hamiltonian.*

At the same time, we find a 2-connected $\{K_{1,3}, P_6\}$-heavy graph which is not Hamiltonian (see Fig. 2).



Fig. 2. A 2-connected $\{K_{1,3}, P_6\}$-heavy non-Hamiltonian graph ($r \geq 5$).

Besides, we can also construct a 2-connected claw-free and P_6-heavy graph which is not Hamiltonian. This can be shown as follows: Let G be the graph in Fig. 2, where $r \geq 15$ is an integer divisible by 3. Let V_1, V_2, V_3 be a balanced partition of K_r and G' be the graph obtained from G by deleting all the edges in $\bigcup_{i=1}^{3}\{x_i v : v \in V_i\}$. Then G' is a 2-connected claw-free and P_6-heavy graph which is not Hamiltonian.

Note that W contains induced P_4, P_5, C_3, Z_1, Z_2 and B. So we have

Theorem 5. *Let R and S be connected graphs with $R, S \neq P_3$ and let G be a 2-connected graph. Then G being $\{R, S\}$-heavy implies G is Hamiltonian if and only if (up to symmetry) $R = K_{1,3}$ and $S = P_4, P_5, C_3, Z_1, Z_2, B, N$ or W.*

Thus, Theorem 5 gives a complete answer to Problem 1.

For claw-heavy graphs, Chen *et al.* get the following result.

Theorem 6 (Chen, Zhang and Qiao [4]). *Let G be a 2-connected graph. If G is claw-heavy and moreover, $\{P_7, D\}$-free or $\{P_7, H\}$-free, then G is Hamiltonian (see Fig. 3).*



Fig. 3. Graphs D and H.

It is clear that every P_6-free graph is also $\{P_7, D\}$-free. Thus we have that every 2-connected claw-heavy and P_6-free graph is Hamiltonian. Together with Theorems 3 and 4, we have the following characterization:

Theorem 7. *Let S be a connected graph with $S \neq P_3$ and let G be a 2-connected claw-heavy graph. Then G being S-free implies G is Hamiltonian if and only if $S = P_4, P_5, P_6, C_3, Z_1, Z_2, B, N$ or W.*

The necessity of this theorem follows from Theorem 1 immediately.

It is known that the only 2-connected $\{K_{1,3}, Z_3\}$-free non-Hamiltonian graphs have 9 vertices (see [6]), hence for $n \geq 10$, every 2-connected $\{K_{1,3}, Z_3\}$-free graph is also Hamiltonian. This leads to the following

Problem 2. Is every 2-connected $\{K_{1,3}, Z_3\}$-heavy graph on $n \geq 10$ vertices Hamiltonian?

Instead of Theorems 3 and 4, we prove the following two stronger results.

Theorem 8. *If G is a 2-connected $\{K_{1,3}, N_{1,1,2}, D\}$-heavy graph, then G is Hamiltonian (see Fig. 4).*

Theorem 9. *If G is a 2-connected $\{K_{1,3}, N_{1,1,2}, H_{1,1}\}$-heavy graph, then G is Hamiltonian (see Fig. 4).*



Fig. 4. Graphs $N_{1,1,2}$ and $H_{1,1}$.

Note that Brousek [3] gave a complete characterization of triples of connected graphs $K_{1,3}, X, Y$ such that a graph G being 2-connected and $\{K_{1,3}, X, Y\}$-free implies G is Hamiltonian. Clearly, if $K_{1,3}, S, T$ is a triple such that every 2-connected $\{K_{1,3}, S, T\}$-heavy graph is Hamiltonian, then, for some triple $K_{1,3}, X, Y$ of [3], S and T are induced subgraphs of X and Y, respectively (of course, the triples of Theorems 8 and 9 have this property). We refer an interested reader to [3] for more details.

2 Some preliminaries

We first give some additional terminology and notation.

Let G be a graph and X be a subset of $V(G)$. The subgraph of G induced by the set X is denoted $G[X]$. We use $G - X$ to denote the subgraph induced by $V(G) \setminus X$.

Throughout this paper, k and ℓ will always denote positive integers, and we use s and t to denote integers which may be nonpositive. For $s \leq t$, we use $[x_s, x_t]$ to denote the set $\{x_s, x_{s+1}, \ldots, x_t\}$. If $[x_s, x_t]$ is a subset of the vertex set of a graph G, we use $G[x_s, x_t]$, instead of $G[[x_s, x_t]]$, to denote the subgraph induced by $[x_s, x_t]$ in G.

For a path P and $x, y \in V(P)$, $P[x, y]$ denotes the subpath of P from x to y. Similarly, for a cycle C with a given orientation and $x, y \in V(C)$, $\overrightarrow{C}[x, y]$ or $\overleftarrow{C}[y, x]$ denotes the (x, y)-path on C traversed in the same or opposite direction with respect to the given orientation of C.

Let G be a graph and $x_1, x_2, y_1, y_2 \in V(G)$ with $x_1 \neq x_2$ and $y_1 \neq y_2$. Then a subgraph Q of G such that Q has exactly 2 components, each of them being an $(\{x_1, x_2\}, \{y_1, y_2\})$-path, is called an (x_1x_2, y_1y_2)-*disjoint path pair*, or briefly an (x_1x_2, y_1y_2)-*pair* in G.

If G is a graph on $n \geq 2$ vertices, $x \in V(G)$, and a graph G' is obtained from G by adding a (new) vertex y and a pair of edges yx, yz, where z is an arbitrary vertex of G,

$z \neq x$, we say that G' is a *1-extension of* G *at* x *to* y. Similarly, if $x_1, x_2 \in V(G)$, $x_1 \neq x_2$, then the graph G' obtained from G by adding two (new) vertices y_1, y_2 and the edges y_1x_1, y_2x_2 and y_1y_2 is called the *2-extension of* G *at* (x_1, x_2) *to* (y_1, y_2).

Let G be a graph and let $u, v, w \in V(G)$ be distinct vertices of G. We say that G is (u, v, w)*-composed* (or briefly *composed*) if G has a spanning subgraph D (called the *carrier* of G) such that there is an ordering $v_{-k}, \ldots, v_0, \ldots, v_\ell$ $(k, \ell \geq 1)$ of $V(D)$ $(=V(G))$ and a sequence of graphs $D_1, \ldots, D_r$ $(r \geq 1)$ such that

(*a*) $u = v_{-k}$, $v = v_0$, $w = v_\ell$,

(*b*) D_1 is a triangle with $V(D_1) = \{v_{-1}, v_0, v_1\}$,

(*c*) $V(D_i) = [v_{-k_i}, v_{\ell_i}]$ for some k_i, ℓ_i, $1 \leq k_i \leq k$, $1 \leq \ell_i \leq \ell$, and D_{i+1} satisfies one of the following:

  (*i*) D_{i+1} is a 1-extension of D_i at v_{-k_i} to v_{-k_i-1} or at v_{ℓ_i} to v_{ℓ_i+1},

  (*ii*) D_{i+1} is a 2-extension of D_i at (v_{-k_i}, v_{ℓ_i}) to $(v_{-k_i-1}, v_{\ell_i+1})$,

$i = 1, \ldots, r-1$,

(*d*) $D_t = D$.

The ordering $v_{-k}, \ldots, v_0, \ldots, v_\ell$ will be called a *canonical ordering* and the sequence $D_1, \ldots, D_r$ a *canonical sequence* of D (and also of G). Note that a composed graph G can have several carriers, canonical orderings and canonical sequences. Clearly, a composed graph G and any its carrier D are 2-connected, for any canonical ordering, $P = v_{-k} \cdots v_0 \cdots v_\ell$ is a Hamilton path in D (called a *canonical path*), and if $D_1, \ldots, D_r$ is a canonical sequence, then any D_i is $(v_{-k_i}, v_0, v_{\ell_i})$-composed, $i = 1, \ldots, r$.

Now we give a lemma on composed graphs which will be needed in our proofs.

Lemma 1. *Let* G *be a composed graph and let* D *and* $v_{-k}, \ldots, v_0, \ldots, v_\ell$ *be a carrier and a canonical ordering of* G*. Then*

(*i*) D *has a Hamilton* (v_0, v_{-k})*-path,*

(*ii*) *for every* $v_s \in V(G) \setminus \{v_{-k}\}$*,* D *has a spanning* (v_0v_ℓ, v_sv_{-k})*-pair.*

Proof. Let $D_1, \ldots, D_r$ be a canonical sequence and Q the canonical path of D corresponding to the given ordering and, for every $s \in [-k, \ell] \setminus \{0\}$, let $\hat{s}$, $1 \leq \hat{s} \leq r$, be the smallest integer for which $v_s \in V(D_{\hat{s}})$. Clearly, $d_{D_{\hat{s}}}(v_s) = 2$.

Now we prove (*i*) by induction on $|V(D)|$. If $|V(D)| = 3$, the assertion is trivially true. Suppose now that $|V(D)| \geq 4$ and the assertion is true for every graph with at most $|V(D)| - 1$ vertices. By the definition of a carrier, we have the following two cases.

Case 1. $V(D_{r-1}) = [v_{-k+1}, v_\ell]$ and D is a 1-extension of D_{r-1} at v_{-k+1} to v_{-k}.

By the induction hypothesis, D_{r-1} has a Hamilton (v_0, v_{-k+1})-path P'. Then $P = v_0 P' v_{-k+1} v_{-k}$ is a Hamilton (v_0, v_{-k})-path in D.

Case 2. $V(D_{r-1}) = [v_{-k}, v_{\ell-1}]$ and D is a 1-extension of D_{r-1} at $v_{\ell-1}$ to v_ℓ, or $V(D_{r-1}) = [v_{-k+1}, v_{\ell-1}]$ and D is a 2-extension of D_{r-1} at $(v_{-k+1}, v_{\ell-1})$ to (v_{-k}, v_ℓ).

In this case, v_ℓ has a neighbor v_s other than $v_{\ell-1}$, where $s \in [-k, \ell - 2]$.

Case 2.1. $s \in [-k, -2]$.

In this case $s+1 \in [-k+1, -1]$. Consider the graph $D' = D_{\widehat{s+1}}$. Let $V(D') = [v_{s+1}, v_t]$, where $t > 0$. By the induction hypothesis, there exists a Hamilton (v_0, v_t)-path P' of D'. Then the path $P = P' Q[v_t, v_\ell] v_\ell v_s Q[v_s, v_{-k}]$ is a Hamilton (v_0, v_{-k})-path of D.

Case 2.2. $s = -1$.

In this case, the path $P = Q[v_0, v_\ell] v_\ell v_{-1} Q[v_{-1}, v_{-k}]$ is a Hamilton (v_0, v_{-k})-path of D.

Case 2.3. $s \in [0, l - 2]$.

In this case $s + 1 \in [1, \ell - 1]$. Consider the graph $D' = D_{\widehat{s+1}}$. Let $V(D') = [v_t, v_{s+1}]$, where $t < 0$ and $d_{D'}(v_{s+1}) = 2$. By the induction hypothesis, there exists a Hamilton (v_0, v_t)-path P' of D', and the edge $v_s v_{s+1}$ is in $E(P')$ by the fact $d_{D'}(v_{s+1}) = 2$. Thus the path $P = P' - v_s v_{s+1} \cup Q[v_{s+1}, v_l] v_l v_s \cup Q[v_t, v_{-k}]$ is a Hamilton (v_0, v_{-k})-path of G.

So the proof of (i) is complete.

Now we prove (ii). We distinguish the following three cases.

Case 1. $s \in [-k + 1, 0]$.

In this case, $s - 1 \in [-k, -1]$. Consider the graph $D' = D_{\widehat{s-1}}$. Let $V(D') = [v_{s-1}, v_t]$, where $t > 0$ and $d_{D'}(v_{s-1}) = 2$. By (i), there exists a Hamilton (v_0, v_t)-path P' of D' and $v_{s-1} v_s \in E(P')$. Thus $R' = P' - v_{s-1} v_s$ is a spanning $(v_0 v_t, v_s v_{s-1})$-pair of D', and $R = R' \cup Q[v_t, v_l] \cup Q[v_{s-1}, v_{-k}]$ is a spanning $(v_0 v_\ell, v_s v_{-k})$-pair of D.

Case 2. $s = 1$.

In this case, $R = Q[v_0, v_{-k}] \cup Q[v_1, v_\ell]$ is a spanning $(v_0 v_\ell, v_1 v_{-k})$-pair of D.

Case 3. $s \in [2, \ell]$.

In this case, $s - 1 \in [1, l - 1]$. Consider the graph $D' = D_{\widehat{s-1}}$. Let $V(D') = [v_t, v_{s-1}]$, where $t < 0$. By (i), there exists a Hamilton (v_0, v_t)-path P' of G'. Thus $P_1 = P' Q[v_t, v_{-k}]$ and $P_2 = Q[v_s, v_\ell]$ form a spanning $(v_0 v_\ell, v_s v_{-k})$-pair of D.

The proof is complete. □

Let G be a graph on n vertices and $k \geq 3$ an integer. A sequence of vertices $C = v_1v_2\cdots v_kv_1$ such that for all $i \in [1,k]$ either $v_iv_{i+1} \in E(G)$ or $d(v_i) + d(v_{i+1}) \geq n$ (indices modulo k) is called an *Ore-cycle* or briefly, *o-cycle* of G. The *deficit* of an o-cycle C is the integer $\mathrm{def}(C) = |\{i \in [1,k] : v_iv_{i+1} \notin E(G)\}|$. Thus, a cycle is an o-cycle of deficit 0. Similarly we define an *o-path* of G.

Now, we prove the following lemma on o-cycles.

Lemma 2. *Let G be a graph and let C' be an o-cycle in G. Then there is a cycle C in G such that $V(C') \subset V(C)$.*

Proof. Let C_1 be an o-cycle in G such that $V(C') \subset V(C_1)$ and $\mathrm{def}(C_1)$ is smallest possible, and suppose, to the contrary, that $\mathrm{def}(C_1) \geq 1$. Without loss of generality suppose that $C_1 = v_1v_2\ldots v_kv_1$, where $v_1v_k \notin E(G)$ and $d(v_1) + d(v_k) \geq n$. We use P to denote the o-path $P = v_1v_2\cdots v_k$.

If v_1 and v_k have a common neighbor $x \in V(G) \setminus V(P)$, then $C_2 = v_1Pv_kxv_1$ is an o-cycle in G with $V(C') \subset V(C_2)$ and $\mathrm{def}(C_2) < \mathrm{def}(C_1)$, a contradiction. Hence $N_{G-P}(v_1) \cap N_{G-P}(v_k) = \emptyset$. Then we have $d_P(v_1) + d_P(v_k) \geq |V(P)|$ since $d(v_1) + d(v_k) \geq n$. Thus, there exists $i \in [2, k-1]$ such that $v_i \in N_P(v_1)$ and $v_{i-1} \in N_P(v_k)$, and then again $C_2 = v_1P[v_1, v_{i-1}]v_{i-1}v_kP[v_k, v_i]v_iv_1$ is an o-cycle with $V(C') \subset V(C_2)$ and $\mathrm{def}(C_2) < \mathrm{def}(C_1)$, a contradiction. $\square$

Note that Lemma 2 immediately implies that if P is an (x,y)-path or an o-path in G with $|V(P)|$ larger than the length of a longest cycle in G, then $xy \notin E(G)$ and $d(x) + d(y) < n$.

In the following, we denote $\overline{E}(G) = \{uv : uv \in E(G) \text{ or } d(u) + d(v) \geq n\}$.

Let C be a cycle in G, $x, x_1, x_2 \in V(C)$ three distinct vertices, and set $X = V(Q)$, where Q is the (x_1, x_2)-path on C containing x. We say that the pair of vertices (x_1, x_2) is *x-good on C*, if for some $j \in \{1, 2\}$ there is a vertex $x' \in X \setminus \{x_j\}$ such that

(1) there is an (x, x_{3-j})-path P such that $V(P) = X \setminus \{x_j\}$,

(2) there is an $(xx_{3-j}, x'x_j)$-pair D such that $V(D) = X$,

(3) $d(x_j) + d(x') \geq n$.

Lemma 3. *Let G be a graph, and C be a cycle of G with a given orientation. Let $x, y \in V(C)$ and let R be an (x,y)-path in G which is internally disjoint with C. If there are vertices $x_1, x_2, y_1, y_2 \in V(C) \setminus \{x, y\}$ such that*

(i) x_2, x, x_1, y_1, y, y_2 appear in this order along $\overrightarrow{C}$ (possibly $x_1 = y_1$ or $x_2 = y_2$),

(ii) (x_1, x_2) *is* x*-good on* C*,*

(iii) (y_1, y_2) *is* y*-good on* C*,*

then there is a cycle C' *in* G *such that* $V(C) \cup V(R) \subset V(C')$*.*

Proof. Assume the opposite. Let P_1 and D_1 be the path and disjoint path pair associated with x, and P_2 and D_2 associated with y; and let $Q_1 = \overrightarrow{C}[x_1, y_1]$ and $Q_2 = \overleftarrow{C}[x_2, y_2]$.

By the definition of an x-good pair, without loss of generality, we can assume that P_1 is an (x, x_1)-path, D_1 is an $(xx_1, x'x_2)$-pair, and $d(x_2) + d(x') \geq n$.

Case 1. P_2 is a (y, y_1)-path, D_2 is a $(yy_1, y'y_2)$-pair, and $d(y_2) + d(y') \geq n$.

In this case the path $P = Q_2 \cup D_2 \cup R \cup P_1 \cup Q_1$ is an (x_2, y')-path which contains all the vertices in $V(C) \cup V(R)$, and $P' = Q_2 \cup D_1 \cup R \cup P_2 \cup Q_1$ is an (x', y_2)-path which contains all the vertices in $V(C) \cup V(R)$. Thus, by Lemma 2, $d(x_2) + d(y') < n$ and $d(x') + d(y_2) < n$, a contradiction to $d(x_2) + d(x') \geq n$ and $d(y_2) + d(y') \geq n$.

Case 2. P_2 is a (y, y_2)-path, D_2 is a $(yy_2, y'y_1)$-pair, and $d(y_1) + d(y') \geq n$.

Case 2.1. The $(xx_1, x'x_2)$-pair D_1 is formed by an (x, x_2)-path and an (x_1, x')-path.

In this case, the path $P = Q_2 \cup P_2 \cup R \cup P_1 \cup Q_1$ is an (x_2, y_1)-path which contains all the vertices in $V(C) \cup V(R)$, and the path $P' = D_1 \cup Q_1 \cup Q_2 \cup R \cup D_2$ is an (x', y')-path which contains all the vertices in $V(C) \cup V(R)$. By Lemma 2, $d(x_2) + d(y_1) < n$ and $d(x') + d(y') < n$, a contradiction.

Case 2.2. The $(xx_1, x'x_2)$-pair D_1 is formed by an (x, x')-path and an (x_1, x_2)-path.

Case 2.2.1. The $(yy_2, y'y_1)$-pair D_2 is formed by an (y, y_1)-path and an (y_2, y')-path.

This case can be proved similarly as in Case 2.1.

Case 2.2.2. The $(yy_2, y'y_1)$-pair D_2 is formed by an (y, y')-path and an (y_1, y_2)-path.

In this case, the path $P = Q_2 \cup D_2 \cup R \cup P_1 \cup Q_1$ is an (x_2, y')-path containing all vertices in $V(C) \cup V(R)$, and the path $P' = Q_2 \cup D_1 \cup R \cup P_2 \cup Q_1$ is an (x', y_1)-path containing all vertices in $V(C) \cup V(R)$. By Lemma 2, $d(x_2) + d(y') < n$ and $d(x') + d(y_1) < n$, a contradiction.

The proof is complete. □

3 Proof of Theorem 8

Let C be a longest cycle of G with a given orientation, set $n = |V(G)|$ and $c = |V(C)|$, and assume that G is not Hamiltonian, i.e. $c < n$. Then $V(G)\backslash V(C) \neq \emptyset$. Since G is 2-connected, there exists a (u_0, v_0)-path with length at least 2 which is internally disjoint from C, where $u_0, v_0 \in V(C)$. Let $R = z_0z_1z_2\cdots z_{r+1}$, where $z_0 = u_0$ and $z_{r+1} = v_0$, be such a path, and choose R as short as possible. Let r_1 and r_2 denote the number of interior vertices in $\overrightarrow{C}[u_0, v_0]$ and $\overrightarrow{C}[v_0, u_0]$, respectively (note that clearly $r_1 + r_2 + 2 = c$). We denote the vertices of C by $\overrightarrow{C} = u_0u_1u_2\cdots u_{r_1}v_0u_{-r_2}u_{-r_2+1}\cdots u_{-1}u_0$ or $\overleftarrow{C} = v_0v_1v_2\cdots v_{r_1}u_0v_{-r_2}v_{-r_2+1}\cdots v_{-1}v_0$, where $u_\ell = v_{r_1+1-\ell}$ and $u_{-k} = v_{-r_2-1+k}$ (see Fig. 5). Let H be the component of $G - C$ which contains the vertices in $[z_1, z_r]$.



Fig. 5. $C \cup R$, the subgraph of G.

Claim 1. Let $x \in V(H)$ and $y \in \{u_1, u_{-1}, v_1, v_{-1}\}$. Then $xy \notin \overline{E}(G)$.

Proof. Without loss of generality, we assume $y = u_1$. Let P' be an (x, z_1)-path in H. Then $P = P'z_1u_0\overleftarrow{C}[u_0, u_1]$ is an (x, y)-path which contains all the vertices in $V(C) \cup V(P')$. By Lemma 2, we have $xy \notin \overline{E}(G)$. □

Claim 2. $u_1u_{-1} \in \overline{E}(G)$, $v_1v_{-1} \in \overline{E}(G)$.

Proof. If $u_1u_{-1} \notin E(G)$, by Claim 1, the graph induced by $\{u_0, z_1, u_1, u_{-1}\}$ is a claw, where $d(z_1) + d(u_{\pm 1}) < n$. Since G is a claw-heavy graph, we have that $d(u_1) + d(u_{-1}) \geq n$.

The second assertion can be proved similarly. □

Claim 3. $u_1v_{-1} \notin \overline{E}(G)$, $u_{-1}v_1 \notin \overline{E}(G)$, $u_0v_{\pm 1} \notin \overline{E}(G), u_{\pm 1}v_0 \notin \overline{E}(G)$.

Proof. Since $P = \overrightarrow{C}[u_1, v_0]R\overleftarrow{C}[u_0, v_{-1}]$ is a (u_1, v_{-1})-path which contains all the vertices in $V(C) \cup V(R)$, we have $u_1v_{-1} \notin \overline{E}(G)$ by Lemma 2.

If $u_0v_1 \in \overline{E}(G)$, then $Pc = \overrightarrow{C}[u_1, v_1]v_1u_0R\overrightarrow{C}[v_0, u_{-1}]u_{-1}u_1$ is an o-cycle which contains all the vertices of $V(C) \cup V(R)$. By Lemma 2, there exists a cycle which contains all the vertices in $V(C) \cup V(R)$, a contradiction.

The other assertions can be proved similarly. $\square$

Claim 4. Either $u_1u_{-1} \in E(G)$ or $v_1v_{-1} \in E(G)$.

Proof. Assume the opposite. By Claim 2 we have $d(u_1)+d(u_{-1}) \geq n$ and $d(v_1)+d(v_{-1}) \geq n$. By Claim 3, we have $d(u_1) + d(v_{-1}) < n$ and $d(u_{-1}) + d(v_1) < n$, a contradiction. $\square$

Now, we distinguish two cases.

Case 1. $r \geq 2$, or $r = 1$ and $u_0v_0 \notin E(G)$.

By Claim 4, without loss of generality, we assume that $u_1u_{-1} \in E(G)$. Thus $G[u_{-1}, u_1]$ is (u_{-1}, u_0, u_1)-composed.

Claim 5. $z_2u_0 \notin \overline{E}(G)$.

Proof. By the choice of the path R, we have $z_2u_0 \notin E(G)$. Now we prove that $d(z_2) + d(u_0) < n$.

Claim 5.1. Every neighbor of u_0 is in $V(C) \cup V(H)$; every neighbor of z_2 is in $V(C) \cup V(H)$.

Proof. Assume the opposite. Let $z' \in V(H')$ be a neighbor of u_0 where H' is a component of $G - C$ other than H. Then we have $z'z_1 \notin E(G)$ and $N_{G-C}(z') \cap N_{G-C}(z_1) = \emptyset$.

By Claim 1, we have $u_1z_1 \notin \overline{E}(G)$, and similarly $u_1z' \notin \overline{E}(G)$. Thus the graph induced by $\{u_0, u_1, z_1, z'\}$ is a claw, where $d(u_1) + d(z_1) < n$ and $d(u_1) + d(z') < n$. Then we have $d(z_1) + d(z') \geq n$.

Since $N_{G-C}(z_1) \cap N_{G-C}(z') = \emptyset$, there exist two vertices $x_1, x_2 \in V(C)$ such that $x_1x_2 \in E(\overrightarrow{C})$ and $z_1x_1, z'x_2 \in E(G)$. Thus $P = z_1x_1\overleftarrow{C}[x_1, x_2]x_2z'$ is a (z_1, z')-path which contains all the vertices in $V(C) \cup \{z_1, z'\}$. By Lemma 2, there exists a cycle which contains all the vertices in $V(C) \cup \{z_1, z'\}$, a contradiction.

If $z_2 = v_0$, the second assertion can be proved similarly; and if $z_2 \neq v_0$, the assertion is obvious. $\square$

Let $h = |V(H)|$ and $k = |N_H(u_0)|$. Then we have $d_H(z_2) + d_H(u_0) \leq h + k$. Since $z_1 \in N_H(u_0)$, we have $k \geq 1$. Let $N_H(u_0) = \{y_1, y_2, \ldots, y_k\}$, where $y_1 = z_1$.

Claim 5.2. $y_iy_j \in \overline{E}(G)$ for all $1 \leq i < j \leq k$.

Proof. If $y_iy_j \notin E(G)$, then by Claim 1, the graph induced by $\{u_0, u_1, y_i, y_j\}$ is a claw, where $d(y_i) + d(u_1) < n$ and $d(y_j) + d(u_1) < n$. Thus we have $d(y_i) + d(y_j) \geq n$. ☐

Now, let Q be the o-path $Q = z_2y_1y_2 \cdots y_ku_0$. It is clear that $R[z_2, v_0]$ and Q are internally disjoint, and Q contains at least k vertices in $V(H)$. In the following, we use C' to denote the cycle $\overrightarrow{C}[u_1, u_{-1}]u_{-1}u_1$ if $z_2 \neq v_0$, and to denote the o-cycle $\overrightarrow{C}[u_1, v_1]v_1v_{-1}\overrightarrow{C}[v_{-1}, u_{-1}]u_{-1}u_1$ if $z_2 = v_0$.

By Claims 1 and 3, we have $z_2v_{r_1} \notin E(G)$, where $v_{r_1} = u_1$. Let v_ℓ be the last vertex in $\overleftarrow{C}[v_1, u_1]$ such that $z_2v_\ell \in E(G)$. If there are no neighbors of z_2 in $\overleftarrow{C}[v_1, u_1]$, then let $v_\ell = v_0$.

Claim 5.3. For every vertex $v_{\ell'} \in N_{[v_1,v_{r_1}]}(z_2) \cup \{v_0\}$, $u_0v_{\ell'+1} \notin E(G)$.

Proof. By Claim 3, we have $u_0v_1 \notin E(G)$.

If $z_2v_{\ell'} \in E(G)$ and $u_0v_{\ell'+1} \in E(G)$, then $C'' = \overrightarrow{C'}[v_{\ell'}, v_{\ell'+1}]v_{\ell'+1}u_0Qz_2v_{\ell'}$ is an o-cycle which contains all the vertices in $V(C) \cup V(Q)$, a contradiction. ☐

Claim 5.4. $r_1 - \ell \geq k + 1$, and for every vertex $v_{\ell'} \in [v_{\ell+1}, v_{\ell+k}]$, $u_0v_{\ell'} \notin E(G)$.

Proof. Assume the opposite. Let $v_{\ell'}$ be the first vertex in $[v_{\ell+1}, v_{r_1}]$ such that $u_0v_{\ell'} \in E(G)$, and $\ell' - \ell < k + 1$.

If $v_\ell = v_0$, then $C'' = \overrightarrow{C}[v_0, u_{-1}]u_{-1}u_1\overrightarrow{C}[u_1, v_{\ell'}]v_{\ell'}u_0QR[z_2, v_0]$ is an o-cycle which contains all the vertices in $V(C)\backslash[v_1, v_{\ell'-1}] \cup V(Q)$, and $|V(C'')| > c$, a contradiction.

Thus, we assume that $v_\ell \neq v_0$, and $z_2v_\ell \in E(G)$. Then $C'' = \overrightarrow{C'}[v_\ell, v_{\ell'}]v_{\ell'}u_0Qz_2v_\ell$ is an o-cycle which contains all the vertices in $V(C)\backslash[v_{\ell+1}, v_{\ell'-1}] \cup V(Q)$, and $|V(C'')| > c$, a contradiction.

Thus we have $\ell' - \ell \geq k + 1$. Note that $u_0v_{r_1} \in E(G)$, we have $r_1 - \ell \geq k + 1$. ☐

Let $d_1 = |N_{[v_1,v_{r_1}]}(z_2) \cup \{v_0\}|$, $d_2 = |N_{[v_{-r_2},v_{-1}]}(z_2) \cup \{v_0\}|$, $d_1' = |N_{[v_1,v_{r_1}]}(u_0)|$ and $d_2' = |N_{[v_{-r_2},v_{-1}]}(u_0)|$. Then $d_C(z_2) \leq d_1 + d_2 - 1$ and $d_C(u_0) \leq d_1' + d_2' + 1$.

By Claims 5.3 and 5.4, we have $d_1' \leq r_1 - d_1 - k + 1$, and similarly, $d_2' \leq r_2 - d_2 - k + 1$. Thus $d_C(z_2) + d_C(u_0) \leq r_1 + r_2 - 2k + 2 = c - 2k$. Note that $d_H(z_2) + d_H(u_0) \leq h + k$. By Claim 5.1, we have $d(z_2) + d(u_0) \leq c + h - k < n$. ☐

Recall that $G[u_{-1}, u_1]$ is (u_{-1}, u_0, u_1)-composed. Now we prove the following claims.

Claim 6. If $G[u_{-k}, u_\ell]$ is (u_{-k}, u_0, u_ℓ)-composed with canonical ordering $u_{-k}, u_{-k+1}, \ldots, u_\ell$, then $k \leq r_2 - 2$ and $\ell \leq r_1 - 2$.

Proof. Let $D_1, D_2, \ldots, D_r$ be a canonical sequence of $G[u_{-k}, u_\ell]$ corresponding to the canonical ordering $u_{-k}, u_{-k+1}, \ldots, u_\ell$. Suppose that $k > r_2 - 2$. Consider the the graph $D' = D_{\widehat{-r_2+1}}$, where $\widehat{-r_2+1}$ be the smallest integer such that $u_{-r_2+1} \in V(D_{\widehat{-r_2+1}})$. Let $V(D') = [u_{-r_2+1}, u_{\ell'}]$. By Lemma 1, there exists a $(u_0, u_{\ell'})$-path P such that $V(P) = [u_{-r_2+1}, u_{\ell'}]$. Then $C' = v_{-1}v_0RP\overrightarrow{C}[u_{\ell'}, v_1]v_1v_{-1}$ is an o-cycle which contains all the vertices in $V(C) \cup V(R)$, a contradiction. $\square$

Claim 7. If $G[u_{-k}, u_\ell]$ is (u_{-k}, u_0, u_ℓ)-composed with canonical ordering $u_{-k}, u_{-k+1}, \ldots, u_\ell$, where $k \leq r_2 - 2$ and $l \leq r_1 - 2$, and any two nonadjacent vertices in $[u_{-k-1}, u_{\ell+1}]$ have degree sum less than n, then one of the following is true:
(1) $G[u_{-k-1}, u_\ell]$ is (u_{-k-1}, u_0, u_ℓ)-composed with canonical ordering $u_{-k-1}, u_{-k}, \ldots, u_\ell$,
(2) $G[u_{-k}, u_{\ell+1}]$ is $(u_{-k}, u_0, u_{\ell+1})$-composed with canonical ordering $u_{-k}, u_{-k+1}, \ldots, u_{\ell+1}$,
(3) $G[u_{-k-1}, u_{\ell+1}]$ is $(u_{-k-1}, u_0, u_{\ell+1})$-composed with canonical ordering $u_{-k-1}, u_{-k}, \ldots, u_{\ell+1}$.

Proof. Assume the opposite, which implies that for every vertex $u_s \in [u_{-k+1}, u_\ell]$, $u_{-k-1}u_s \notin E(G)$, and for every vertex $u_s \in [u_{-k}, u_{\ell-1}]$, we have $u_{\ell+1}u_s \notin E(G)$ and $u_{-k-1}u_{\ell+1} \notin E(G)$.

Claim 7.1. Let $z \in \{z_1, z_2\}$ and $u_s \in [u_{-k-1}, u_{\ell+1}] \backslash \{u_0\}$. Then $zu_s \notin \overline{E}(G)$.

Proof. Without loss of generality, we assume that $s > 0$. If $s = 1$, the assertion is true by Claims 1 and 3. So we assume that $s \in [2, \ell+1]$ and $s - 1 \in [1, \ell]$. By the definition of a composed graph, there exists $t \in [-k, -1]$ such that $G[u_t, u_{s-1}]$ is (u_t, u_0, u_{s-1})-composed. By Lemma 1, there exists a (u_0, u_t)-path P' such that $V(P') = [u_t, u_{s-1}]$.

If $z \neq v_0$, then $P = R[z, u_0]P'\overleftarrow{C}[u_t, u_s]$ is a (z, u_s)-path which contains all the vertices in $V(C) \cup \{z\}$. By Lemma 2, we have $zu_s \notin \overline{E}(G)$.

If $z = v_0$ and $v_0u_s \in \overline{E}(G)$, then $C' = RP'\overleftarrow{C}[u_t, v_{-1}]v_{-1}v_1\overleftarrow{C}[v_1, u_s]u_sv_0$ is an o-cycle which contains all the vertices in $V(C) \cup V(R)$, a contradiction. $\square$

Let $G' = G[[u_{-k-1}, u_\ell] \cup \{z_1, z_2\}]$ and $G'' = G[[u_{-k-1}, u_{\ell+1}] \cup \{z_1, z_2\}]$.

Claim 7.2. G'', and then G', is $\{K_{1,3}, N_{1,1,2}\}$-free.

Proof. By Claims 5 and 7.1, and the condition that any two nonadjacent vertices in $[u_{-k-1}, u_{\ell+1}]$ have degree sum less than n, we have that any two nonadjacent vertices in G'' have degree sum less than n. Since G (and then G'') is $\{K_{1,3}, N_{1,1,2}\}$-heavy, we have that G'' is $\{K_{1,3}, N_{1,1,2}\}$-free. $\square$

Claim 7.3. $N_{G'}(u_0) \backslash \{z_1\}$ is a clique.

Proof. If there are two vertices $x, x' \in N_{G'}(u_0)\backslash\{z_1\}$ such that $xx' \notin E(G')$, then the graph induced by $\{u_0, z_1, x, x'\}$ is a claw, a contradiction. $\square$

Now, we define $N_i = \{x \in V(G') : d_{G'}(x, u_{-k-1}) = i\}$. Then we have $N_0 = \{u_{-k-1}\}$, $N_1 = \{u_{-k}\}$ and $N_2 = N_{G'}(u_{-k})\backslash\{u_{-k-1}\}$.

By the definition of a composed graph, we have $|N_2| \geq 2$. If there are two vertices $x, x' \in N_2$ such that $xx' \notin E(G')$, then the graph induced by $\{u_{-k}, u_{-k-1}, x, x'\}$ is a claw, a contradiction. Thus, N_2 is a clique.

We assume $u_0 \in N_j$, where $j \geq 2$. Then $z_1 \in N_{j+1}$ and $z_2 \in N_{j+2}$.

If $|N_i| = 1$ for some $i \in [2, j-1]$, say, $N_i = \{x\}$, then x is a cut vertex of the graph $G[u_{-k}, u_l]$. By the definition of a composed graph, $G[u_{-k}, u_l]$ is 2-connected. This implies $|N_i| \geq 2$ for every $i \in [2, j-1]$.

Claim 7.4. For $i \in [1, j]$, N_i is a clique.

Proof. We prove this claim by induction on i. For $i = 1, 2$, the claim is true by the analysis above. So we assume that $3 \leq i \leq j$, and we have that $N_{i-3}, N_{i-2}, N_{i-1}, N_{i+1}$ and N_{i+2} is nonempty, and $|N_{i-1}| \geq 2$.

First we choose a vertex $x \in N_i$ which has a neighbor $y \in N_{i+1}$ such that it has a neighbor $z \in N_{i+2}$. We prove that for every $x' \in N_i$, $xx' \in E(G)$. We assume that $xx' \notin E(G)$.

If $x'y \in E(G)$, then the graph induced by $\{y, x, x', z\}$ is a claw, a contradiction. Thus, we have $x'y \notin E(G)$. If x and x' have a common neighbor in N_{i-1}, denote it by w, then let v be a neighbor of w in N_{i-2}, and the graph induced by $\{w, v, x, x'\}$ is a claw, a contradiction. Thus we have that x and x' have no common neighbors in N_{i-1}.

Let w be a neighbor of x in N_{i-1} and w' be a neighbor of x' in N_{i-1}. Then $xw', x'w \notin E(G)$. Let v be a neighbor of w in N_{i-2} and u be a neighbor of v in N_{i-3}. If $w'v \notin E(G)$, then the graph induced by $\{w, v, w', x\}$ is a claw, a contradiction. Thus we have $w'v \in E(G)$, and then the graph induced by $\{v, u, w', x', w, x, y\}$ is an $N_{1,1,2}$, a contradiction.

Thus we have $xx' \in E(G)$ for every $x' \in N_i$.

Now, let x' and x'' be two vertices in N_i other than x such that $x'x'' \notin E(G)$. We have $xx', xx'' \in E(G)$.

If $x'y \in E(G)$, then similarly to the case of x, we have $x'x'' \in E(G)$, a contradiction. Thus we have $x'y \notin E(G)$. Similarly, $x''y \notin E(G)$. Then the graph induced by $\{x, x', x'', y\}$ is a claw, a contradiction.

Thus, N_i is a clique. $\square$

If there exists some vertex $y \in N_{j+1}$ other than z_1, then we have $yu_0 \notin E(G)$ by Claim 7.3. Let x be a neighbor of y in N_j, w be a neighbor of u_0 in N_{j-1} and v be a neighbor of w in N_{j-2}. Then $xu_0 \in E(G)$ by Claim 7.4 and $xw \in E(G)$ by Claim 7.3. Thus the graph induced by $\{w, v, x, y, u_0, z_1, z_2\}$ is an $N_{1,1,2}$, a contradiction. So we assume that all vertices in $[u_{-k}, u_\ell]$ are in $\bigcup_{i=1}^{j} N_i$.

If $u_\ell \in N_j$, then let w be a neighbor of u_0 in N_{j-1} and v be a neighbor of w in N_{j-2}. Then the graph induced by $\{w, v, u_0, z_1, u_\ell, u_{\ell+1}\}$ is an $N_{1,1,2}$, a contradiction. Thus we have that $u_\ell \notin N_j$ and then $j \geq 3$.

Let $u_\ell \in N_i$, where $i \in [2, j-1]$. If u_ℓ has a neighbor in N_{i+1}, then let y be a neighbor of u_ℓ in N_{i+1}, and w be a neighbor of u_ℓ in N_{i-1}. Then the graph induced by $\{u_\ell, w, y, u_{\ell+1}\}$ is a claw, a contradiction. So we have that u_ℓ has no neighbors in N_{i+1}.

Let $x \in N_i$ be a vertex other than u_ℓ which has a neighbor y in N_{i+1} such that it has a neighbor z in N_{i+2}. Let w be a neighbor of x in N_{i-1}, and v be a neighbor of w in N_{i-2}. If $u_\ell w \notin E(G)$, then the graph induced by $\{x, w, u_\ell, y\}$ is a claw, a contradiction. So we have that $u_\ell w \in E(G)$. Then the graph induced by $\{w, v, u_\ell, u_{\ell+1}, x, y, z\}$ is an $N_{1,1,2}$, a contradiction.

Thus the claim holds. ☐

Now we choose k, ℓ such that

(1) $G[u_{-k}, u_\ell]$ is (u_{-k}, u_0, u_ℓ)-composed with canonical ordering $u_{-k}, u_{-k+1}, \dots, u_\ell$;
(2) any two nonadjacent vertices in $[u_{-k}, u_\ell]$ have degree sum less than n; and
(3) $k + \ell$ is as big as possible.

By Claim 7, we have that there exists a vertex $u_s \in [u_{-k+1}, u_\ell]$ such that $d(u_{-k-1}) + d(u_s) \geq n$, or there exists a vertex $u_s \in [u_{-k}, u_{\ell-1}]$ such that $d(u_s) + d(u_{\ell+1}) \geq n$, or $d(u_{-k-1}) + d(u_{\ell+1}) \geq n$. Thus, we have

Claim 8. (u_{-k-1}, u_ℓ) or $(u_{-k}, u_{\ell+1})$ or $(u_{-k-1}, u_{\ell+1})$ is u_0-good on C.

Proof. If there exists a vertex $u_s \in [u_{-k+1}, u_\ell]$ such that $d(u_{-k-1}) + d(u_s) \geq n$, then, by Lemma 1, there exists a (u_0, u_ℓ)-path P such that $V(P) = [u_{-k}, u_\ell]$, and there exists a $(u_0 u_\ell, u_s u_{-k})$-pair D' such that $V(D') = [u_{-k}, u_\ell]$, and $D = D' + u_{-k} u_{-k-1}$ is a $(u_0 u_\ell, u_s u_{-k-1})$-pair such that $V(D) = [u_{-k-1}, u_\ell]$. Thus (u_{-k-1}, u_ℓ) is u_0-good on C.

If there exists a vertex $u_s \in [u_{-k}, u_{\ell-1}]$ such that $d(u_s) + d(u_{\ell+1}) \geq n$, we can prove the result similarly.

If $d(u_{-k-1}) + d(u_{\ell+1}) \geq n$, then by Lemma 1, there exists a (u_0, u_ℓ)-path P' such that $V(P') = [u_{-k}, u_\ell]$ and there exists a (u_0, u_{-k})-path P'' such that $V(P'') = [u_{-k}, u_\ell]$. Then

$P = P'u_1u_{\ell+1}$ is a $(u_0, u_{\ell+1})$-path such that $V(P) = [u_{-k}, u_{\ell+1}]$, and $D = P''u_{-k}u_{-k-1} \cup u_{\ell+1}$ is a $(u_0u_{\ell+1}, u_{\ell+1}u_{-k-1})$-pair such that $V(D) = [u_{-k-1}, u_{\ell+1}]$. Thus $(u_{-k-1}, u_{\ell+1})$ is u_0-good on C. $\square$

Claim 9. There exist $v_{-k'} \in V(\overrightarrow{C}[v_{-1}, u_{-k-1}])$ and $v_{\ell'} \in V(\overleftarrow{C}[v_1, u_{\ell+1}])$ such that $(v_{-k'}, v_{\ell'})$ is is v_0-good on C.

Proof. By Claim 6, we have $k \leq r_2 - 2$ and $l \leq r_1 - 2$.

If $v_1v_{-1} \notin E(G)$, then by Claim 2, $d(v_1)+d(v_{-1}) \geq n$. Then $P = v_0v_1$ is a (v_0, v_1)-path and $D = v_0v_{-1} \cup v_1$ is a $(v_0v_1, v_{-1}v_1)$-pair. Thus we have that (v_{-1}, v_1) is v_0-good on C.

Now we assume that $v_1v_{-1} \in E(G)$, and then $G[v_{-1}, v_1]$ is (v_{-1}, v_0, v_1)-composed.

Let $r_2' = r_2 - k$ and $r_1' = r_1 - \ell$.

Claim 9.1. If $G[v_{-k'}, v_{\ell'}]$ is $(v_{-k'}, v_0, v_{\ell'})$-composed with canonical ordering $v_{-k'}, v_{-k'+1}, \ldots, v_{\ell'}$, then $k' \leq r_2' - 1$ and $\ell' \leq r_1' - 1$.

Proof. Let $D_1, D_2, \ldots, D_r$ be a canonical sequence of $G[v_{-k'}, v_{\ell'}]$ corresponding to the canonical ordering $v_{-k'}, v_{-k'+1}, \ldots, v_{\ell'}$. Suppose that $k' > r_2' - 1$. Consider the the graph $D' = D_{\widehat{-r_2'}}$, where $\widehat{-r_2'}$ is the smallest integer such that $v_{-r_2'} \in V(D_{\widehat{-r_2'}})$. Let $V(D') = [v_{-r_2'}, v_{\ell''}]$. By Lemma 1, there exists a $(v_0, v_{\ell''})$-path P such that $V(P) = [v_{-r_2'}, u_{\ell''}]$. Then $C' = P\overrightarrow{C}[u_\ell, v_{\ell''}]P'R$ is a cycle which contains all the vertices in $V(C) \cup V(R)$, a contradiction. $\square$

Similarly to Claim 7, we have

Claim 9.2. If $G[v_{-k'}, v_{\ell'}]$ is $(v_{-k'}, v_0, v_{\ell'})$-composed with canonical ordering $v_{-k'}, v_{-k'+1}, \ldots, v_{\ell'}$, where $k' \leq r_2' - 1$ and $\ell \leq r_1' - 1$, and any two nonadjacent vertices in $[v_{-k'-1}, v_{\ell'+1}]$ have degree sum less than n, then one of the following is true:

(1) $G[v_{-k'-1}, v_{\ell'}]$ is $(v_{-k'-1}, v_0, v_{\ell'})$-composed with canonical ordering $v_{-k'-1}, v_{-k'}, \ldots, v_{\ell'}$,

(2) $G[v_{-k'}, v_{l'+1}]$ is $(v_{-k'}, v_0, v_{\ell'+1})$-composed with canonical ordering $v_{-k'}, v_{-k'+1}, \ldots, v_{\ell'+1}$,

(3) $G[v_{-k'-1}, v_{l'+1}]$ is $(v_{-k'-1}, v_0, v_{\ell'+1})$-composed with canonical ordering $v_{-k'-1}, v_{-k'}, \ldots, v_{\ell'+1}$.

Now we choose k', ℓ' such that

(1) $G[v_{-k'}, v_{\ell'}]$ is $(v_{-k'}, v_0, v_{\ell'})$-composed with canonical ordering $v_{-k'}, v_{-k'+1}, \ldots, v_{\ell'}$;

(2) any two nonadjacent vertices in $[v_{-k'}, v_{\ell'}]$ have degree sum less than n; and

(3) $k' + \ell'$ is as big as possible.

Similarly to Claim 8, we have $(v_{-k'-1}, v_{l'})$ or $(v_{-k'}, v_{l'+1})$ or $(v_{-k'-1}, v_{l'+1})$ is v_0-good on C. $\square$

From Claims 8 and 9, we get that there exists a cycle which contains all the vertices in $V(C) \cup V(R)$ by Lemma 3, a contradiction.

Case 2. $r = 1$ and $u_0v_0 \in E(G)$.

We have $u_0u_{-1} \in E(G)$ and $u_0u_{-r_2} \notin E(G)$, where $u_{-r_2} = v_{-1}$. Let u_{-k-1} be the first vertex in $\overleftarrow{C}[u_{-1}, v_{-1}]$ such that $u_0u_{-k-1} \notin E(G)$. Then $k \leq r_2 - 1$.

Similarly, let $v_{\ell+1}$ be the first vertex in $\overleftarrow{C}[v_1, u_1]$ such that $v_0v_{\ell+1} \notin E(G)$. Then $\ell \leq r_1 - 1$.

Claim 10. Let $x \in [u_{-k-1}, u_{-1}]$ and $y \in [v_1, v_{\ell+1}]$. Then

(i) $xz_1, xv_0 \notin \overline{E}(G)$,

(ii) $yz_1, yu_0 \notin \overline{E}(G)$,

(iii) $xy \notin \overline{E}(G)$.

Proof. *(i)* If $x = u_{-1}$, then by Claims 1 and 3, we have $u_{-1}z_1, u_{-1}v_0 \notin \overline{E}(G)$. So we assume that $x = u_{-k'}$ where $-k' \in [-k-1, -2]$ and $u_0u_{-k'+1} \in E(G)$.

If $u_{-k'}z_1 \in \overline{E}(G)$, then $C' = u_0u_{-k'+1}\overrightarrow{C}[u_{-k'+1}, u_{-1}]u_{-1}u_1\overrightarrow{C}[u_1, u_{-k'}]u_{-k'}z_1u_0$ is an o-cycle which contains all the vertices in $V(C) \cup V(R)$, a contradiction.

If $u_{-k'}v_0 \in \overline{E}(G)$, then $C' = u_0u_{-k'+1}\overrightarrow{C}[u_{-k'+1}, u_{-1}]u_{-1}u_1\overrightarrow{C}[u_1, v_1]v_1v_{-1}\overrightarrow{C}[v_{-1}, u_{-k'}]$ $u_{-k'}v_0R$ is an o-cycle which contains all the vertices in $V(C) \cup V(R)$, a contradiction.

The assertion *(ii)* can be proved similarly.

(iii) If $x = u_{-1}$ and $y = v_1$, then by Claim 3, we have $xy \notin \overline{E}(G)$.

If $u_{-k'}v_1 \in \overline{E}(G)$, where $k' \in [2, k+1]$, then $C' = u_0R\overrightarrow{C}[v_0, u_{-k'}]u_{-k'}v_1\overleftarrow{C}[v_1, u_1]u_1u_{-1}$ $\overleftarrow{C}[u_{-1}, u_{-k'+1}]u_{-k'+1}u_0$ is an o-cycle which contains all the vertices in $V(C) \cup V(R)$, a contradiction.

If $u_{-1}v_{\ell'} \in \overline{E}(G)$, where $\ell' \in [2, \ell+1]$, then we can prove the result similarly.

If $u_{-k'}v_{\ell'} \in \overline{E}(G)$, where $k' \in [2, k+1]$ and $\ell' \in [2, \ell+1]$, then $C' = u_0u_{-k'+1}\overrightarrow{C}[u_{-k'+1}, u_{-1}]$ $u_{-1}u_1\overrightarrow{C}[u_1, v_{l'}]v_{l'}u_{-k'}\overleftarrow{C}[u_{-k'}, v_{-1}]v_{-1}v_1\overleftarrow{C}[v_1, v_{l'-1}]v_{l'-1}v_0R$ is an o-cycle which contains all the vertices in $V(C) \cup V(R)$, a contradiction. ☐

Claim 11. Either $u_{-k-1}u_0 \notin \overline{E}(G)$, or $v_{\ell+1}v_0 \notin \overline{E}(G)$.

Proof. Assume the opposite. Since $u_{-k-1}u_0, v_{\ell+1}v_0 \notin E(G)$, we have $d(u_{-k-1}) + d(u_0) \geq n$ and $d(v_{\ell+1}) + d(v_0) \geq n$. By Claim 10, we have $d(u_0) + d(v_{\ell+1}) < n$ and $d(v_0) + d(u_{-k-1}) < n$, a contradiction. ☐

Without loss of generality, we assume that $u_{-k-1}u_0 \notin \overline{E}(G)$. If $v_{\ell+1}v_0 \notin \overline{E}(G)$, then the subgraph induced by $\{z_1, v_0, v_\ell, v_{\ell+1}, u_0, u_{-k}, u_{-k-1}\}$ is a D which is not heavy, a contradiction. Since $v_0v_{\ell+1} \notin E(G)$, we have $d(v_0) + d(v_{\ell+1}) \geq n$.

Claim 12. Either (v_{-1}, v_1) or $(v_{-1}, v_{\ell+1})$ is v_0-good on C.

Proof. If $v_1v_{-1} \notin E(G)$, then, by Claim 2, $d(v_1) + d(v_{-1}) \geq n$. Then $P = v_0v_1$ is a (v_0, v_1)-path and $D = v_0v_{-1} \cup v_1$ is a $(v_0v_1, v_{-1}v_1)$-pair. Thus, (v_{-1}, v_1) is v_0-good on C.

If $v_1v_{-1} \in E(G)$, then $P = v_0v_\ell\overrightarrow{C}[v_\ell, v_1]v_1v_{-1}$ is a (v_0, v_{-1})-path and $D = v_0 \cup v_{-1}v_1\overleftarrow{C}[v_1, v_{\ell+1}]$ is a (v_0v_{-1}, v_0v_{l+1})-pair. Since $d(v_0) + d(v_{\ell+1}) \geq n$, we have that $(v_{-1}, v_{\ell+1})$ is v_0-good on C. ☐

Claim 13. If $G[u_{-k'}, u_{\ell'}]$ is $(u_{-k'}, u_0, u_{\ell'})$-composed with canonical ordering $u_{-k'}, u_{-k'+1}, \ldots, u_{\ell'}$, then $k' \leq r_2 - 2$ and $\ell' \leq r_1 - \ell - 1$.

Proof. The claim can be proved similarly as Claims 6 and 9.1. ☐

Now we prove the following claim.

Claim 14. If $G[u_{-k'}, u_{\ell'}]$ is $(u_{-k'}, u_0, u_{\ell'})$-composed with canonical ordering $u_{-k'}, u_{-k'+1}, \ldots, u_{\ell'}$, where $k' \leq r_2 - 2$ and $\ell' \leq r_1 - \ell - 1$, and any two nonadjacent vertices in $[u_{-k'-1}, u_{\ell'+1}]$ have degree sum less than n, then one of the following is true:

(1) $G[u_{-k'-1}, u_{\ell'}]$ is $(u_{-k'-1}, u_0, u_{\ell'})$-composed with canonical ordering $u_{-k'-1}, u_{-k'}, \ldots, u_{\ell'}$,

(2) $G[u_{-k'}, u_{\ell'+1}]$ is $(u_{-k'}, u_0, u_{\ell'+1})$-composed with canonical ordering $u_{-k'}, u_{-k'+1}, \ldots, u_{\ell'+1}$,

(3) $G[u_{-k'-1}, u_{\ell'+1}]$ is $(u_{-k'-1}, u_0, u_{\ell'+1})$-composed with canonical ordering $u_{-k'-1}, u_{-k'}, \ldots, u_{\ell'+1}$.

Proof. Assume the opposite, which implies that for every vertex $u_s \in [u_{-k'+1}, u_{\ell'}]$, $u_{-k'-1}u_s \notin E(G)$, and for every vertex $u_s \in [u_{-k'}, u_{\ell'-1}]$, we have $u_{\ell'+1}u_s \notin E(G)$ and $u_{-k'-1}u_{\ell'+1} \notin E(G)$.

Claim 14.1. Let $v \in \{v_0, v_1\}$ and $u_s \in [u_{-k'-1}, u_{\ell'+1}]\backslash\{u_0\}$. Then $vu_s \notin \overline{E}(G)$.

Proof. Similarly to Claim 7.1, we have $v_0u_s \notin \overline{E}(G)$.

Now we assume that $v_1u_s \in \overline{E}(G)$.

Note that if $v_0v_2 \notin E(G)$, then $d(v_0) + d(v_2) \geq n$. We have $v_0v_2 \in \overline{E}(G)$.

If $s \in [-k' - 1, -2]$, then $s + 1 \in [-k', -1]$. By the definition of a composed graph, there exists $t \in [1, \ell']$ such that $G[u_{s+1}, u_t]$ is (u_{s+1}, u_0, u_t)-composed. By Lemma 1, there exists a (u_0, u_t)-path P such that $V(P) = [u_{s+1}, u_t]$. Then $C' = P\overrightarrow{C}[u_t, v_1]v_1u_s\overleftarrow{C}[u_s, v_0]R$ is an o-cycle which contains all the vertices in $V(C) \cup V(R)$, a contradiction.

If $s = -1$, by Claim 3, we have $v_1u_{-1} \notin \overline{E}(G)$.

If $s=1$, then $C'=\overleftarrow{C}[u_0,v_{-1}]v_{-1}v_1u_1\overrightarrow{C}[u_1,v_2]v_2v_0R$ is an o-cycle which contains all the vertices in $V(C)\cup V(R)$, a contradiction.

If $s\in[2,\ell'+1]$, then $s-1\in[1,\ell']$. By the definition of a composed graph, there exists $t\in[-k',-1]$ such that $G[u_t,u_{s-1}]$ is (u_t,u_0,u_{s-1})-composed. By Lemma 1, there exists a (u_0,u_t)-path P such that $V(P)=[u_t,u_{s-1}]$. Then $C'=P\overleftarrow{C}[u_t,v_{-1}]v_{-1}v_1u_s\overrightarrow{C}[u_s,v_2]v_2v_0R$ is an o-cycle which contains all the vertices in $V(C)\cup V(R)$, a contradiction. $\square$

Let $G'=G[[u_{-k'-1},u_{\ell'}]\cup\{v_0,v_1\}]$ and $G''=G[[u_{-k'-1},u_{\ell'+1}]\cup\{v_0,v_1\}]$. Then, similarly to Claim 7.2, we have

Claim 14.2. G'', and then G', is $\{K_{1,3},N_{1,1,2}\}$-free.

Thus we can prove the claim similarly to Claim 7. $\square$

Now we choose k',ℓ' such that

(1) $G[u_{-k'},u_{\ell'}]$ is $(u_{-k'},u_0,u_{\ell'})$-composed;

(2) any two nonadjacent vertices in $[u_{-k'},u_{\ell'}]$ have degree sum less than n; and

(3) $k'+\ell'$ is as big as possible.

Similarly to Claim 8, we have

Claim 15. $(u_{-k'-1},u_{\ell'})$ or $(u_{-k'},u_{\ell'+1})$ or $(u_{-k'-1},u_{\ell'+1})$ is u_0-good on C.

By Claim 13, we have $k'\le r_2-2$ and $\ell'\le r_1-\ell-2$.

From Claims 12 and 15, we can get that there exists a cycle which contains all vertices in $V(C)\cup V(R)$ by Lemma 3, a contradiction.

The proof is complete.

4 Proof of Theorem 9

Let C be a longest cycle of G with a given orientation. We use n to denote the order of G, and c the length of C. Assume that G is not Hamiltonian. Then $V(G)\backslash V(C)\neq\emptyset$. Since G is 2-connected, there exists a (u_0,v_0)-path of length at least 2 which is internally disjoint with C, where $u_0,v_0\in V(C)$. Let $R=z_0z_1z_2\cdots z_{r+1}$ be such a path which is as short as possible, where $z_0=u_0$ and $z_{r+1}=v_0$. We assume that the length of $\overrightarrow{C}[u_0,v_0]$ is r_1+1 and the length of $\overrightarrow{C}[v_0,u_0]$ is r_2+1, where $r_1+r_2+2=c$. We denote the vertices of C by $\overrightarrow{C}=u_0u_1u_2\cdots u_{r_1}v_0u_{-r_2}u_{-r_2+1}\cdots u_{-1}u_0$ or $\overleftarrow{C}=v_0v_1v_2\cdots v_{r_1}u_0v_{-r_2}v_{-r_2+1}\cdots v_{-1}v_0$, where $u_\ell=v_{r_1+1-\ell}$ and $u_{-k}=v_{-r_2-1+k}$. Let H be the component of $G-C$ which contains the vertices in $[z_1,z_r]$.

As in Section 3, we have the following claims.

Claim 1. Let $x \in V(H)$ and $y \in \{v_1, v_{-1}, u_1, u_{-1}\}$. Then $xy \notin \overline{E}(G)$.

Claim 2. $u_1u_{-1} \in \overline{E}(G)$, $v_1v_{-1} \in \overline{E}(G)$.

Claim 3. $u_1v_{-1} \notin \overline{E}(G)$, $u_{-1}v_1 \notin \overline{E}(G)$, $u_0v_{\pm1} \notin \overline{E}(G), u_{\pm1}v_0 \notin \overline{E}(G)$.

Claim 4. Either u_1u_{-1} or v_1v_{-1} is in $E(G)$.

By Claim 4, without loss of generality, we assume that $u_1u_{-1} \in E(G)$. Then we have that $G[u_{-1}, u_1]$ is (u_{-1}, u_0, u_1)-composed.

Claim 5. If $G[u_{-k}, u_\ell]$ is (u_{-k}, u_0, u_ℓ)-composed, then $k \leq r_2 - 2$ and $\ell \leq r_1 - 2$.

The proof of Claim 5 is similar to that of Claim 6 in Section 3.

Now we prove the following claim.

Claim 6. If $G[u_{-k}, u_\ell]$ is (u_{-k}, u_0, u_ℓ)-composed with canonical ordering $u_{-k}, u_{-k+1}, \dots, u_\ell$, where $k \leq r_2 - 2$ and $l \leq r_1 - 2$, and any two nonadjacent vertices in $[u_{-k-1}, u_{\ell+1}]$ have degree sum less than n, then one of the following is true:
(1) $G[u_{-k-1}, u_\ell]$ is (u_{-k-1}, u_0, u_ℓ)-composed with canonical ordering $u_{-k-1}, u_{-k}, \dots, u_\ell$,
(2) $G[u_{-k}, u_{\ell+1}]$ is $(u_{-k}, u_0, u_{\ell+1})$-composed with canonical ordering $u_{-k}, u_{-k+1}, \dots, u_{\ell+1}$,
(3) $G[u_{-k-1}, u_{\ell+1}]$ is $(u_{-k-1}, u_0, u_{\ell+1})$-composed with canonical ordering $u_{-k-1}, u_{-k}, \dots, u_{\ell+1}$.

Proof. Assume the opposite, which implies that for every vertex $u_s \in [u_{-k+1}, u_\ell]$, $u_{-k-1}u_s \notin E(G)$, and for every vertex $u_s \in [u_{-k}, u_{\ell-1}]$, $u_{\ell+1}u_s \notin E(G)$ and $u_{-k-1}u_{\ell+1} \notin E(G)$.

Claim 6.1. For every vertex $z \in \{z_1, z_2\}$ and $u_s \in [u_{-k-1}, u_{\ell+1}] \backslash \{u_0\}$ we have $zu_s \notin \overline{E}(G)$; and if $z_2u_0 \notin E(G)$, then also $z_2u_0 \notin \overline{E}(G)$.

This claim can be proved similarly to Claims 5 and 7.1 in Section 3.

Let $G' = G[[u_{-k-1}, u_\ell] \cup \{z_1, z_2\}]$ and $G'' = G[[u_{-k-1}, u_{\ell+1}] \cup \{z_1, z_2\}]$.

Similarly to Claims 7.2 and 7.3 in Section 3, we have

Claim 6.2. G'', and then G', is $\{K_{1,3}, N_{1,1,2}, H_{1,1}\}$-free.

Claim 6.3. $N_{G'}(u_0) \backslash \{z_1, z_2\}$ is a clique.

Now, we define $N_i = \{x \in V(G') : d_{G'}(x, u_{-k-1}) = i\}$. Then we have $N_0 = \{u_{-k-1}\}$, $N_1 = \{u_{-k}\}$ and $N_2 = N_{G'}(u_{-k}) \backslash \{u_{-k-1}\}$.

By the definition of a composed graph, we have that $|N_2| \geq 2$. If there are two vertices $x, x' \in N_2$ such that $xx' \notin E(G')$, then the graph induced by $\{u_{-k}, u_{-k-1}, x, x'\}$ is a claw. Thus N_2 is a clique.

We assume $u_0 \in N_j$, where $j \geq 2$. Then $z_1 \in N_{j+1}$; and $z_2 \in N_{j+1}$ if $z_2u_0 \in E(G)$ and $z_2 \in N_{j+2}$ if $z_2u_0 \notin E(G)$.

If $|N_i| = 1$ for some $i \in [2, j-1]$, say, $N_i = \{x\}$, then x is a cut vertex of the graph $G[u_{-k}, u_\ell]$. By the definition of a composed graph, $G[u_{-k}, u_\ell]$ is 2-connected. This implies $|N_i| \geq 2$ for every $i \in [2, j-1]$.

Claim 6.4. For $i \in [1, j]$, N_i is a clique.

Proof. If $i < j$, or $i = j$ and $z_2u_0 \notin E(G)$, then we can prove the assertion similarly to Claim 7.4 in Section 3. Thus we assume that $i = j$ and $z_2u_0 \in E(G)$.

If $j = 2$, the assertion is true by the analysis above. So we assume that $j \geq 3$, and we have that $N_{j-3}, N_{j-2}, N_{j-1}, N_{j+1}$ is nonempty and $|N_{j-1}| \geq 2$.

First we prove that for every $x \in N_j \backslash \{u_0\}$, $u_0x \in E(G)$. We assume that $u_0x \notin E(G)$.

By Claim 6.1 we have $xz_1 \notin E(G)$. If u_0 and x have a common neighbor in N_{j-1}, denoted w, then let v be a neighbor of w in N_{j-2}; but then the graph induced by $\{w, v, u_0, x\}$ is a claw, a contradiction. Thus we have that u_0 and x have no common neighbors in N_{j-1}.

Let w be a neighbor of u_0 in N_{j-1} and w' be a neighbor of x in N_{j-1}. Then $u_0w', xw \notin E(G)$. Let v be a neighbor of w in N_{j-2} and u be a neighbor of v in N_{j-3}. If $w'v \notin E(G)$, then the graph induced by $\{w, v, w', u_0\}$ is a claw, a contradiction. Thus we have $w'v \in E(G)$, and then the graph induced by $\{v, u, w', x, w, u_0, z_1\}$ is an $N_{1,1,2}$, a contradiction.

Thus we have $u_0x \in E(G)$ for every $x \in N_j$. Then by Claim 6.3, we have that N_j is a clique. □

If there exists some vertex $y \in N_{j+1}$ other than z_1 and z_2, then we have $yu_0 \notin E(G)$ by Claim 6.3. Let x be a neighbor of y in N_j, w be a neighbor of u_0 in N_{j-1} and v be a neighbor of w in N_{j-2}. Then $xu_0 \in E(G)$ by Claim 6.4 and $xw \in E(G)$ by Claim 6.3. Thus the graph induced by $\{w, v, x, y, u_0, z_1, z_2\}$ is an $N_{1,1,2}$ if $z_2u_0 \notin E(G)$, and is an $H_{1,1}$ if $z_2u_0 \in E(G)$, a contradiction. So we assume that all vertices in $[u_{-k}, u_\ell]$ are in $\bigcup_{i=1}^{j} N_i$.

If $u_\ell \in N_j$, then let w be a neighbor of u_0 in N_{j-1} and v be a neighbor of w in N_{j-2}. Then the graph induced by $\{w, v, u_0, z_1, u_\ell, u_{\ell+1}\}$ is an $N_{1,1,2}$ if $z_2u_0 \notin E(G)$, and is an $H_{1,1}$ if $z_2u_0 \in E(G)$, a contradiction. Thus we have that $u_\ell \notin N_j$ and then $j \geq 3$.

Let $u_\ell \in N_i$, where $i \in [2, j-1]$. If u_ℓ has a neighbor in N_{i+1}, then let y be a neighbor of u_ℓ in N_{i+1}, and w be a neighbor of u_ℓ in N_{i-1}. Then the graph induced by $\{u_\ell, w, y, u_{\ell+1}\}$ is a claw, a contradiction. Thus we have that u_ℓ has no neighbors in N_{i+1}.

Let $x \in N_i$ be a vertex other than u_ℓ which has a neighbor y in N_{i+1} such that it has a neighbor z in N_{i+2}. Let w be a neighbor of x in N_{i-1}, and v be a neighbor of w in N_{i-2}.

If $u_\ell w \notin E(G)$, then the graph induced by $\{x, w, u_\ell, y\}$ is a claw, a contradiction. Thus we have that that $u_\ell w \in E(G)$. Then the graph induced by $\{w, v, u_\ell, u_{\ell+1}, x, y, z\}$ is an $N_{1,1,2}$, a contradiction.

Thus the claim holds. $\square$

Now we choose k, ℓ such that

(1) $G[u_{-k}, u_\ell]$ is (u_{-k}, u_0, u_ℓ)-composed with canonical ordering $u_{-k}, u_{-k+1}, \ldots, u_\ell$;

(2) any two nonadjacent vertices in $[u_{-k}, u_\ell]$ have degree sum less than n; and

(3) $k + \ell$ is as big as possible.

Similarly to Claims 8 and 9 in Section 3, we have

Claim 7. (u_{-k-1}, u_ℓ) or $(u_{-k}, u_{\ell+1})$ or $(u_{-k-1}, u_{\ell+1})$ is u_0-good on C.

Claim 8. There exist $v_{-k'} \in V(\overrightarrow{C}[v_{-1}, u_{-k-1}])$ and $v_{\ell'} \in V(\overleftarrow{C}[v_1, u_{\ell+1}])$ such that $(v_{-k'}, v_{\ell'})$ is v_0-good on C.

From Claims 7 and 8, we can get that there exists a cycle which contains all the vertices in $V(C) \cup V(R)$ by Lemma 3, a contradiction.

The proof is complete.

References

[1] P. Bedrossian, Forbidden subgraph and minimum degree conditons for hamiltonicity, Ph.D Thesis, Memphis State University, USA, 1991.

[2] J.A. Bondy and U.S.R. Murty, Graph Theory with Applications, Macmillan London and Elsevier, New York (1976).

[3] J. Brousek, Forbidden triples for hamiltonicity, Discrete Math. 251 (2002) 71-76.

[4] B. Chen, S. Zhang and S. Qiao, Hamilton cycles in claw-heavy graphs, Discrete Math. 309 (2009) 2015-2019.

[5] G. Fan, New sufficient conditions for cycles in graphs, J. Combin. Theory B 37 (1984) 221-227.

[6] R.J. Faudree, R.J. Gould, Z. Ryjáček and I. Schiermeyer, Forbidden subgraphs and cycle extendability, J. Combin. Math. Combin. Comput. 19 (1995) 109-128.